

Euler Sums of Hyperharmonic Numbers

Ayhan Dil

Department of Mathematics,
Akdeniz University, 07058 Antalya Turkey
adil@akdeniz.edu.tr

Khristo N. Boyadzhiev

Department of Mathematics and Statistics,
Ohio Northern University Ada, Ohio 45810, USA,
k-boyadzhiev@onu.edu

September 5, 2012

Abstract

The hyperharmonic numbers $h_n^{(r)}$ are defined by means of the classical harmonic numbers. We show that the Euler-type sums with hyperharmonic numbers:

$$\sigma(r, m) = \sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n^m}$$

can be expressed in terms of series of Hurwitz zeta function values. This is a generalization of a result of Mezo and Dil. We also provide an explicit evaluation of $\sigma(r, m)$ in a closed form in terms of zeta values and Stirling numbers of the first kind. Furthermore, we evaluate several other series involving hyperharmonic numbers.

2000 Mathematics Subject Classification 11B73, 11M99.

Key words: Riemann zeta function, Hurwitz zeta function, Euler sums, harmonic and hyperharmonic numbers, Stirling numbers, Beta function.

1 Introduction

In this paper we are interested in Euler-type sums with hyperharmonic numbers. Such series could be of interest in analytic number theory. We will show that these sums are related to the values of the Riemann zeta function. In [15] the authors considered the case $r = 1$. Here we extend this result to $r > 1$.

In the second section we express $\sigma(r, m)$ as a special series involving zeta values. In the third section we evaluate $\sigma(r, m)$ as a finite sum including Stirling numbers of the first kind, zeta values, and values of the digamma (psi) function.

In the last fourth section we use certain integral representaions to evaluate several series with hyperharmonic numbers. For example,

$$\sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n(n+1)\dots(n+r)} = \frac{\pi^2}{6r!},$$

$$\sum_{n=1}^{\infty} h_n^{(r)} B(r+1, n+1) = 1$$

and

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{(n+1)} h_n^{(2)} = \sum_{n=1}^{\infty} (-1)^n H_n \beta(n+1).$$

1.1 Definitions and notation

Harmonic and Hyperharmonic numbers

The n -th harmonic number is defined by the n -th partial sum of the harmonic series as

$$H_n := \sum_{k=1}^n \frac{1}{k}, \quad (1)$$

where $H_0 = 0$.

For an integer $r > 1$, let

$$h_n^{(r)} := \sum_{k=1}^n h_k^{(r-1)} \quad (2)$$

with $h_n^{(1)} := H_n$, be the n -th hyperharmonic number of order r ([9]).

These numbers can be expressed in terms of binomial coefficients and ordinary harmonic numbers as ([9, 15]):

$$h_n^{(r)} = \binom{n+r-1}{r-1} (H_{n+r-1} - H_{r-1}). \quad (3)$$

The well-known generating functions of the harmonic and hyperharmonic numbers are given as

$$\sum_{n=1}^{\infty} H_n x^n = -\frac{\ln(1-x)}{1-x} \quad (4)$$

and

$$\sum_{n=1}^{\infty} h_n^{(r)} x^n = -\frac{\ln(1-x)}{(1-x)^r} \quad (5)$$

respectively.

Euler discovered the following formula:

$$2 \sum_{n=1}^{\infty} \frac{H_n}{n^m} = (m+2) \zeta(m+1) - \sum_{n=1}^{m-2} \zeta(m-n) \zeta(n+1), \quad m = 2, 3, \dots \quad (6)$$

where $\zeta(s)$ is the Riemann zeta function ([11, 18]).

For certain pairs of positive integers r and m several authors have evaluated in closed form the Euler sums

$$S(r, m) = \sum_{n=1}^{\infty} \frac{H_n^{(r)}}{n^m} \text{ where } H_n^{(r)} = 1 + \frac{1}{2^r} + \dots + \frac{1}{n^r}$$

(see [4, 11] and for an elementary procedure [6]). In this paper we want to do the same for the sums

$$\sigma(r, m) = \sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n^m}.$$

Considering hyperharmonic generalization of Euler sums, Mezo and Dil ([15]) obtained:

$$\sigma(r, m) = \sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n^m} < \infty \text{ where } m > r$$

Rearrangements of these series gives:

$$\sigma(r, m) = \sum_{n=1}^{\infty} h_n^{(r-1)} \zeta(m, n) \quad (7)$$

where $m > r$ ([15]). Here $\zeta(s, a)$ is the Hurwitz zeta function defined as

$$\zeta(s, a) = \sum_{n=0}^{\infty} \frac{1}{(n+a)^s}$$

where $a > 0$ is a fixed number and $\text{Re}(s) > 1$.

Using (6) and (7) for $m = 2, 3, \dots$ we have

$$2 \sum_{k=1}^{\infty} \frac{\zeta(m, k)}{k} = (m+2) \zeta(m+1) - \sum_{n=1}^{m-2} \zeta(m-n) \zeta(n+1) \quad (8)$$

Hence as a special case of (7) with $r = 1$, Mezo and Dil ([15]) found the following results with the help of (8)

$$\sum_{k=1}^{\infty} \frac{\zeta(2, k)}{k} = 2\zeta(3),$$

$$\sum_{k=1}^{\infty} \frac{\zeta(3, k)}{k} = \frac{5}{4}\zeta(4),$$

$$\begin{aligned}\sum_{k=1}^{\infty} \frac{\zeta(4, k)}{k} &= 3\zeta(5) - \frac{\pi^2}{6}\zeta(3), \\ \sum_{k=1}^{\infty} \frac{\zeta(5, k)}{k} &= \frac{\pi^6}{540} - \frac{1}{2}\zeta^2(3), \\ &\dots\end{aligned}$$

2 Euler Sums of Hyperharmonic Numbers

The following theorem provides a general version of the equation (7).

Theorem 1 *For integers $0 \leq k < r < m$ we have,*

$$\sigma(r, m) = \sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n^m} = \sum_{n=1}^{\infty} h_n^{(r-k-1)} \sum_{n \leq i_1 \leq i_2 \leq \dots \leq i_k \leq \infty} \zeta(m, i_k) \quad (9)$$

where

$$\sum_{n \leq i_1 \leq i_2 \leq \dots \leq i_k \leq \infty} \zeta(m, i_k) = \sum_{i_1=n}^{\infty} \sum_{i_2=i_1}^{\infty} \dots \sum_{i_k=i_{k-1}}^{\infty} \zeta(m, i_k). \quad (10)$$

Proof. From (7) we have

$$\sigma(r, m) = \sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n^m} = \sum_{n=1}^{\infty} \sum_{j=1}^n h_j^{(r-2)} \zeta(m, n). \quad (11)$$

Rearrangements of the RHS of (11) gives

$$\sigma(r, m) = \sum_{n=1}^{\infty} h_n^{(r-2)} \sum_{i_1=n}^{\infty} \zeta(m, i_1). \quad (12)$$

Using a similar argument, after k -steps we obtain the desired result. ■

Remark 2 *Let us consider two special cases of (9). To state Euler sums of hyperharmonic numbers $\sigma(r, m)$ in terms of the multiple sums of Hurwitz zeta function we choose $k = r - 1$ i.e.,*

$$\sigma(r, m) = \sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n^m} = \sum_{n=1}^{\infty} \frac{\sum_{n \leq i_1 \leq i_2 \leq \dots \leq i_{r-1} \leq \infty} \zeta(m, i_{r-1})}{n}. \quad (13)$$

Also the case $k = r - 2$ gives

$$\sigma(r, m) = \sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n^m} = \sum_{n=1}^{\infty} H_n \sum_{n \leq i_1 \leq i_2 \leq \dots \leq i_{r-2} \leq \infty} \zeta(m, i_{r-2}) \quad (14)$$

which is a representation of $\sigma(r, m)$ in terms of harmonic numbers and the multiple sums of Hurwitz zeta function.

3 A closed form of $\sigma(r, m)$

We are interested in Euler sums of hyperharmonic numbers in the form

$$\sigma(r, m) = \sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n^m} \text{ where } m \geq r + 1.$$

In the following theorem we use the series

$$\mu(m, r) = \sum_{n=1}^{\infty} \frac{1}{n^m (n+r)}$$

and the unsigned Stirling numbers of the first kind $\begin{bmatrix} n \\ k \end{bmatrix}$

To obtain a closed form of $\sigma(r, m)$ we need the following lemma.

Lemma 3 (see [5]) *For every positive integer m and every $r > 0$*

$$\mu(m; r) = \sum_{k=1}^{m-1} \frac{(-1)^{k-1} \zeta(m-k+1)}{r^k} + \frac{(-1)^{m-1}}{r^m} (\Psi(r+1) + \gamma). \quad (15)$$

Here $\Psi(s)$ is the digamma function such that $\gamma = -\Psi(1)$ is Euler's constant.

Theorem 4 *For the integers $0 < r < m$ we have*

$$\begin{aligned} \sigma(r, m) &= \frac{1}{(r-1)!} \sum_{k=1}^r \begin{bmatrix} r \\ k \end{bmatrix} \times \\ &\quad \left\{ \zeta_H(m-k+1) - H_{r-1} \zeta(m-k+1) + \sum_{j=1}^{r-1} \mu(m-k+1, j) \right\}. \end{aligned} \quad (16)$$

Proof. Considering the following relation between Stirling numbers of the first kind and the rising factorial,

$$x(x+1) \dots (x+n-1) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix} x^k$$

we get

$$\binom{n+r}{r} = \frac{1}{r!} \sum_{k=1}^{r+1} \begin{bmatrix} r+1 \\ k \end{bmatrix} n^{k-1}.$$

Hence,

$$\begin{aligned} h_n^{(r+1)} &= \frac{1}{r!} \sum_{k=1}^{r+1} \begin{bmatrix} r+1 \\ k \end{bmatrix} n^{k-1} (H_{n+r} - H_r) \\ &= \frac{1}{r!} \sum_{k=1}^{r+1} \begin{bmatrix} r+1 \\ k \end{bmatrix} n^{k-1} \left(H_n + \frac{1}{n+1} + \dots + \frac{1}{n+r} - H_r \right). \end{aligned}$$

Now we can write

$$\frac{h_n^{(r)}}{n^m} = \frac{1}{(r-1)!} \sum_{k=1}^r \begin{bmatrix} r \\ k \end{bmatrix} \left(\frac{H_n}{n^{m-k+1}} - \frac{H_{r-1}}{n^{m-k+1}} + \sum_{j=1}^{r-1} \frac{1}{n^{m-k+1} (n+j)} \right).$$

Therefore,

$$\sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n^m} = \frac{1}{(r-1)!} \sum_{k=1}^r \begin{bmatrix} r \\ k \end{bmatrix} \left\{ \sum_{n=1}^{\infty} \frac{H_n}{n^{m-k+1}} - H_{r-1} \sum_{n=1}^{\infty} \frac{1}{n^{m-k+1}} + \sum_{j=1}^{r-1} \sum_{n=1}^{\infty} \frac{1}{n^{m-k+1} (n+j)} \right\}.$$

In view of (15) we obtain (16). ■

Hereafter we consider the series $\sigma(r, m) = \sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n^m}$ when $m > r > 1$. Using Theorem 4 we obtain representation of the function $\sigma(r, m)$ in terms of the Riemann zeta function and

$$\zeta_H(m) = \sum_{n=1}^{\infty} \frac{H_n}{n^m}.$$

Here are some examples.

Case $r = 2$

By considering the case $r = 2$ in (16) we get

$$\sigma(2, m) = \zeta_H(m-1) + \zeta_H(m) - \zeta(m-1). \quad (17)$$

With the aid of Theorem 1, equation (17) can be written as

$$\sigma(2, m) = \sum_{n=1}^{\infty} H_n \zeta(m, n) = \zeta_H(m-1) + \zeta_H(m) - \zeta(m-1). \quad (18)$$

Setting $m = 3$ in (18) we obtain

$$\sigma(2, 3) = \sum_{n=1}^{\infty} H_n \zeta(3, n) = 2\zeta(3) + \frac{5}{4}\zeta(4) - \zeta(2) \quad (19)$$

and for $m = 4$ we have

$$\sigma(2, 4) = \sum_{n=1}^{\infty} H_n \zeta(4, n) = \frac{5}{4}\zeta(4) + 3\zeta(5) - \zeta(2)\zeta(3) - \zeta(3) \quad (20)$$

and so on.

Case $r = 3$

By (9) and (16) we have

$$\begin{aligned} \sigma(3, m) &= \sum_{n=1}^{\infty} h_n^{(2)} \zeta(m, n) \\ &= \frac{1}{2}\zeta_H(m-2) + \frac{3}{2}\zeta_H(m-1) + \zeta_H(m) - \frac{5}{4}\zeta(m-1) - \frac{3}{4}\zeta(m-2) \end{aligned}$$

Setting $m = 4$ in the above equation we obtain

$$\sigma(3, 4) = \sum_{n=1}^{\infty} h_n^{(2)} \zeta(4, n) = \frac{15}{6} \zeta(4) + \zeta_H(4) - \frac{1}{4} \zeta(3) - \frac{3}{4} \zeta(2). \quad (21)$$

4 Evaluation of some specific hyperharmonic number series

In this section we evaluate some specific series involving hyperharmonic numbers.

Proposition 5

$$\sum_{n=0}^{\infty} \frac{h_n^{(r)}}{(n+1)(n+2)\dots(n+r+1)} = \frac{1}{r!} \quad (22)$$

Proof. Using the formula (see [12])

$$\frac{1}{r!} \int_0^1 t^n (1-t)^r dt = \frac{1}{(n+1)(n+2)\dots(n+r+1)} \quad (23)$$

we can write

$$\frac{1}{r!} \int_0^1 h_n^{(r)} t^n (1-t)^r dt = \frac{h_n^{(r)}}{(n+1)(n+2)\dots(n+r+1)} \quad (24)$$

with the help of (5) we get

$$\sum_{n=0}^{\infty} \frac{h_n^{(r)}}{(n+1)(n+2)\dots(n+r+1)} = -\frac{1}{r!} \int_0^1 \ln(1-t) dt \quad (25)$$

This equation completes the proof since

$$\int_0^1 \ln(1-t) dt = -1. \quad (26)$$

■

Proposition 6

$$\sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n(n+1)\dots(n+r)} = \frac{\pi^2}{6r!} \quad (27)$$

Proof. In the same way as in the above proposition, from (23) we can write

$$\frac{1}{r!} \int_0^1 h_n^{(r)} t^n (1-t)^r \frac{dt}{t} = \frac{h_n^{(r)}}{n(n+1)\dots(n+r)} \quad (28)$$

from which it follows that

$$\sum_{n=1}^{\infty} \frac{h_n^{(r)}}{n(n+1)\dots(n+r)} = -\frac{1}{r!} \int_0^1 \frac{\ln(1-t)}{t} dt. \quad (29)$$

Since (see [12])

$$\int_0^1 \frac{\ln(1-t)}{t} dt = -\frac{\pi^2}{6} \quad (30)$$

we obtain (27). ■

Proposition 7 *We have*

$$\sum_{n=1}^{\infty} \frac{(-1)^n H_n}{n+1} = -\frac{(\ln 2)^2}{2}$$

and

$$\sum_{n=1}^{\infty} \frac{(-1)^{n+1} H_n}{n} = \frac{\pi^2}{12} - \frac{(\ln 2)^2}{2}.$$

Proof. Using the formulas (see [16])

$$\int_{-1}^0 \frac{\ln(1-t)}{1-t} dt = -\frac{(\ln 2)^2}{2}$$

and

$$\int_0^1 \frac{\ln(1+t)}{t(1+t)} dt = \frac{\pi^2}{12} - \frac{(\ln 2)^2}{2}$$

respectively, and considering (4) we obtain the desired results. ■

Proposition 8 *We have*

$$\sum_{n=1}^{\infty} \frac{2(-1)^n (n+2)}{(n+3)(n+4)} h_n^{(4)} = \frac{23}{72} - \frac{(\ln 2)^2}{3}$$

and

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{(n+1)} h_n^{(r+1)} = \frac{2^r - 1}{2^r r^2} - \frac{\ln 2}{2^r r}.$$

Proof. Using the formulas (see [16])

$$\int_0^1 \frac{\ln(1+t)}{(1+t)^4} (1+t^2) dt = \frac{23}{72} - \frac{(\ln 2)^2}{3}$$

and

$$\int_0^1 \frac{\ln(1+t)}{(1+t)^{r+1}} dt = \frac{2^r - 1}{2^r r^2} - \frac{\ln 2}{2^r r}$$

respectively, and considering (5) gives the desired results. ■

Proposition 9 *We have*

$$\sum_{m=1}^{\infty} \frac{(-1)^{m+1} (2m+2n+3)}{(m+1)(m+2n+2)} H_m = 2 \ln 2 \sum_{k=0}^n \frac{1}{2k+1} - \sum_{j=1}^{2n+1} \frac{1}{j} \sum_{k=1}^j \frac{(-1)^{k-1}}{k}$$

and

$$\sum_{m=1}^{\infty} \frac{(-1)^{m+1} 2n}{(m+1)(m+2n+1)} H_m = 2 \ln 2 \sum_{k=0}^{n-1} \frac{1}{2k+1} - \sum_{j=1}^{2n} \frac{1}{j} \sum_{k=1}^j \frac{(-1)^{k-1}}{k}.$$

Proof. Using the formulas (see [16])

$$\int_0^1 \frac{\ln(1+t)}{(1+t)} (1+t^{2n+1}) dt = 2 \ln 2 \sum_{k=0}^n \frac{1}{2k+1} - \sum_{j=1}^{2n+1} \frac{1}{j} \sum_{k=1}^j \frac{(-1)^{k-1}}{k}$$

and

$$\int_0^1 \frac{\ln(1+t)}{(1+t)} (1-t^{2n}) dt = 2 \ln 2 \sum_{k=0}^{n-1} \frac{1}{2k+1} - \sum_{j=1}^{2n} \frac{1}{j} \sum_{k=1}^j \frac{(-1)^{k-1}}{k}$$

respectively completes the proof in view of (4) . ■

Remark 10 *The Beta function $B(x, y)$ is defined as follows (see [17]):*

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt \quad (31)$$

where $\operatorname{Re}(x) > 0$, $\operatorname{Re}(y) > 0$.

Proposition 11 *We have*

$$\sum_{n=1}^{\infty} h_n^{(r)} B(r+1, n+1) = 1 \quad (32)$$

Proof. Using the formula (see [17])

$$\int_0^1 t^n (1-t)^r dt = B(r+1, n+1) \quad (33)$$

and considering (5) gives

$$\sum_{n=1}^{\infty} h_n^{(r)} B(r+1, n+1) = -1 \int_0^1 \ln(1-t) dt$$

This equation completes the proof since (see [12])

$$\int_0^1 \ln(1-t) dt = -1. \quad (34)$$

■

Proposition 12 *We have*

$$\sum_{n=1}^{\infty} h_n^{(r)} B(r+1, n) = \frac{\pi^2}{6}$$

Proof. From equation (33) we have

$$\int_0^1 h_n^{(r)} t^n (1-t)^r \frac{dt}{t} = h_n^{(r)} B(r+1, n).$$

Using (5) this becomes

$$\sum_{n=1}^{\infty} h_n^{(r)} B(r+1, n) = - \int_0^1 \frac{\ln(1-t)}{t} dt.$$

In view of equation (30) we complete the proof. ■

Proposition 13

$$\sum_{n=1}^{\infty} \frac{(-1)^n}{(n+1)} h_n^{(2)} = \sum_{n=1}^{\infty} (-1)^n H_n \beta(n+1)$$

where (see [17])

$$\beta(x) = \frac{1}{2} \left[\Psi\left(\frac{x+1}{2}\right) - \Psi\left(\frac{x}{2}\right) \right] \text{ and } \Psi(x) = \frac{d}{dx} \ln \Gamma(x).$$

Proof. Using the formula ([17])

$$\beta(x) = \int_0^1 \frac{t^{x-1}}{1+t} dt \text{ where } \operatorname{Re}(x) > 0$$

gives

$$\sum_{n=1}^{\infty} (-1)^n H_n \beta(n+1) = \int_0^1 \frac{1}{t+1} \sum_{n=1}^{\infty} (-1)^n H_n t^n dt$$

and now (5) helps to finish the proof. ■

References

- [1] T. M. Apostol and T. H. Vu, Dirichlet series related to the Riemann zeta function. J. Number Theory 19, 85-102 (1982).
- [2] A. T. Benjamin, D. Gaebler and R. Gaebler, A combinatorial approach to hyperharmonic numbers. Integers: Electron. J. Combin. Number Theory 3: 1-9 #A15 (2003).

- [3] D. Borwein and J. M. Borwein, On an intriguing integral and some series related to $\zeta(4)$. Proc. Amer. Math. Soc. 123: 1191–1198 (1995).
- [4] D. Borwein, J. M. Borwein and R. Girgensohn. Explicit evaluation of Euler sums. Proc. Edinburgh Math. Soc. (2) 38 no. 2, 277–294 (1995).
- [5] K. N. Boyadzhiev, Evaluation of Euler-Zagier sums, Int. J. Math. Math. Sci. 27 (2001), no. 7, 407–412.
- [6] K. N. Boyadzhiev, Consecutive Evaluation of Euler Sums. Internat. J. Math. Math. Sci., 29:9, 555–561 (2002).
- [7] K. N. Boyadzhiev, Evaluation of some simple Euler-type series. Far East Journal of Mathematical Sciences (FJMS), 29,(1), 1 -11 (2008).
- [8] Yu. A. Brychkov, O. I. Marichev, A. P. Prudnikov, Tables of Indefinite Integrals, Gordon and Breach, New York-London, 1989.
- [9] J. H. Conway and R. K. Guy, The Book of Numbers. New York, Springer-Verlag, 1996.
- [10] A. Dil and I. Mezo, A Symmetric Algorithm for Hyperharmonic and Fibonacci Numbers. Applied Mathematics and Computation 206, 942–951 (2008).
- [11] P. Flajolet and B. Salvy, Euler Sums and Contour Integral Representations. Experimental Mathematics, 7, 1, (1998).
- [12] A. Jeffrey and D. Zwillinger (editors), Gradshteyn and Ryzhik's Table of Integrals, Series, and Products, Seventh Edition, (2007).
- [13] K. S. Kölbig, Closed Expressions for $\int_0^1 t^{-1} \log^{n-1} t \log^p(1-t) dt$. Mathematics of Computation, Vol. 39, No. 160, pp. 647–654 (1982).
- [14] K. S. Kölbig, Nielsen's generalized polylogarithms, SIAM J. Math. Anal. 17 (1986) 1232–1258.
- [15] I. Mezo and A. Dil, Hyperharmonic series involving Hurwitz zeta function. Journal of Number Theory, 130, 2: 360–369 (2010).
- [16] A. P. Prudnikov, Yu A. Brychkov and O. I. Marichev, Integrals and Series. Elementary Functions. Gordon and Breach, New York-London, 1986.
- [17] A. P. Prudnikov, Yu A. Brychkov and O. I. Marichev, Integrals and Series. Vol. 2. Special Functions. Gordon and Breach, New York-London, 1986.
- [18] R. S. R. C. Rao and A. S. R. Sarma, Some identities involving the Riemann zeta function. Indian J. Pure Appl. Math. 10, 602–607 (1979).
- [19] R. Sitaramachandrarao, A formula of S. Ramanujan. Journal of Number Theory, 25, 1–19 (1987).